

Back-to-back isolated photon-quarkonium production at the LHC and the transverse-momentum-dependent distributions of the gluons in the proton

J.P. Lansberg

IPNO, Université Paris-Sud, CNRS/IN2P3, F-91406, Orsay, France

Jean-Philippe.Lansberg@in2p3.fr

The study of a quasi back-to-back isolated pair made of a heavy quarkonium, such as a J/ψ or a Υ , and a photon produced in proton-proton collisions at the LHC, is probably the optimal way to get right away a first experimental determination of two gluon transverse-momentum-dependent distributions (TMDs) in an unpolarized proton, f_1^g and $h_1^{\perp g}$, the latter giving the distribution of linearly polarized gluons. To substantiate this, we calculate the transverse-momentum-dependent effects that arise in the process under study and discuss the feasibility of their measurements.

Keywords: TMDs, quarkonium, LHC

1. Introduction

It is nowadays well recognized that an important class of observables at particle colliders are transverse^a-momentum distributions of final state events. While a transverse-momentum (q_T) distribution can be described within the standard collinear factorization approach for large q_T , it is convenient and sometimes necessary to resort to an alternative factorization approach for small q_T , typically $q_T \sim M_p \ll Q$, M_p being the proton mass and Q the hard scale of the process under investigation. This alternative approach, called Transverse-Momentum-Dependent (TMD) factorization (see e.g. Ref. 1), takes into account the transverse motion of partons w.r.t. the direction of the parent proton. In this picture the small final-state transverse momentum q_T is a consequence of the parton transverse momenta. Hence, the non-perturbative distribution functions entering a TMD factorization formula not only depend on the collinear momentum fractions x , but also on the transverse momentum p_T .

The information on the transverse-momentum dependence of unpolarized and linearly polarized gluons in an unpolarized proton is encapsulated in the following TMD correlator Φ_g , which describes the transition from a proton to a gluon, such that

$$\begin{aligned} \Phi_g^{\mu\nu}(x, \mathbf{p}_T, \zeta, \mu) &\equiv 2 \int \frac{d(\xi \cdot P) d^2 \xi_T}{(xP \cdot n)^2 (2\pi)^3} e^{i(xP + p_T) \cdot \xi} \text{Tr}_c \left[\langle P | F^{n\nu}(0) \mathcal{U}_{[0, \xi]}^{n[-]} F^{n\mu}(\xi) \mathcal{U}_{[\xi, 0]}^{n[-]} | P \rangle \right]_{\xi \cdot P' = 0} \\ &= -\frac{1}{2x} \left\{ g_T^{\mu\nu} f_1^g - \left(\frac{p_T^\mu p_T^\nu}{M_p^2} + g_T^{\mu\nu} \frac{\mathbf{p}_T^2}{2M_p^2} \right) h_1^{\perp g} \right\}. \end{aligned} \quad (1)$$

^awith respect to the beam axis

The gluon four-momentum p is decomposed as $p = xP + p_T + p^-n$, where n is a lightlike dimensionless vector, conjugated to the momentum P of the parent proton, with no transverse components and satisfying the relation $\zeta^2 = (2n \cdot P)^2/n^2$ (one assumes an analogous decomposition for k and one takes $p^- = k^+ = 0$), $p_T^2 = -\mathbf{p}_T^2$ and $g_T^{\mu\nu} = g^{\mu\nu} - P^\mu n^\nu / P \cdot n - n^\mu P^\nu / P \cdot n$. In addition, $F^{\mu\nu}(x)$ is the gluon field strength. At leading twist, Φ_g is parametrized⁷ in terms of the two gluon TMD distributions discussed above, f_1^g and $h_1^{\perp g}$. In Eq. (1), the gauge link $\mathcal{U}_{[0,\xi]}^{n[-]}$ is needed to render the matrix element gauge invariant. It runs from 0 to ξ via minus infinity along the n direction.

A model-independent positivity bound for $h_1^{\perp g}$ was derived in Ref. 7 and reads

$$\frac{\mathbf{p}_T^2}{2M_p^2} |h_1^{\perp g}(x, \mathbf{p}_T^2)| \leq f_1^g(x, \mathbf{p}_T^2). \quad (2)$$

Since it is T -even, $h_1^{\perp g}$ does not necessarily vanish in absence of initial and final state interactions. This does not prevent it to be nonuniversal if it receives contributions from such interactions.

A number of suggestions to measure these unknown TMDs, f_1^g and $h_1^{\perp g}$, have been discussed in the literature. Whereas $h_1^{\perp g}$ can, in principle, be extracted from the azimuthal dependence of the dijet production in pp collisions⁴, TMD-factorization breaking effects may be significant in this case due to the existence of both initial and final state interactions⁵. To avoid these, one may prefer to rely on heavy-quark or dijet *electro*-production^{2,3} which would however only be studied at a future EIC facility. Diphoton production⁶ should also not be sensitive to factorization-breaking effects. It however suffers from a huge π^0 background and from contaminations from quark-induced channels at RHIC energies which may preclude a clean gluon TMD extraction. It has recently been proposed¹⁰ to look at $C = +1$ quarkonium production in the region where their transverse momentum is smaller than their mass. A one-loop analysis has shown that 1S_0 production may be safer¹³ than that of 3P_J where color-octet contributions might spoil factorization¹³. Low- q_T $C = +1$ quarkonium production may however be very challenging at the LHC; a first study of η_c production has just been carried out by the LHCb but for $q_T > 6$ GeV. Prospects are certainly more promising with the proposed LHC fixed-target experiment AFTER@LHC (see e.g. Ref. 14, 15, 16, 17).

We claim here that polarized and unpolarized gluon TMDs can be accessed right now at the LHC through the reaction¹⁸

$$p(P_A) + p(P_B) \rightarrow Q(P_Q) + \gamma(P_\gamma) + X, \quad (3)$$

where now Q is one of the $C = -1$ charmonium or bottomonium (e.g. J/ψ or Υ) produced almost back-to-back with a photon. In this case, the momentum imbalance of the pair in the final state, $\mathbf{q}_T = \mathbf{P}_{QT} + \mathbf{P}_{\gamma T}$, is small, but not the individual transverse momenta of the two particles. These should be well detectable by the ATLAS or CMS detectors for instance. As we discussed in Ref. 18, both these final states are always produced by gluon fusion (this is in fact also true at lower energies¹⁶), as illustrated by the thin and thicker curves on Fig. 4. In this particular configuration, the process $\Upsilon + \gamma$ is expected to be dominated by the color-singlet contributions¹⁹ (see the solid blue and dashed orange curves on Fig. 4), hence TMD factorization should be applicable since the final-state interactions are expected to be suppressed. In the case of $J/\psi + \gamma$, a precise assesement of a possible color-singlet dominance depends a more precise knowledge of color octet NRQCD matrix elements, which is still lacking^{20,21,22}. To be on the safe side and to single out the color-singlet contribution, it could be useful to isolate the J/ψ . It may also be useful to measure $J/\psi + \gamma$ production (see Ref. 23 and references therein) in general in order to check the possible contribution from Double-Parton Scatterings (DPS).




Fig. 1. Color-octet and color-singlet contributions from $g-g$ fusion and $q-\bar{q}$ annihilation channels to the production of a photon with a) an Υ and b) a J/ψ as a function of Q , the invariant mass of the pair. The curves for the $q-\bar{q}$ annihilation are respectively rescaled by a factor 100 (50) in a) (b)).

It has however recently been emphasised in an analysis of Higgs plus jet production at the LHC²⁴ that it may not be necessary to restrict the analyses to final states made only of colorless particles for the TMD factorisation to be applicable, provided that one of the particles used to measure the momentum imbalance is colorless. If this is the case, color-octet contributions to quarkonium + photon would not break the TMD factorisation. By extension, the study of the momentum imbalance in J/ψ plus jet could also be used to extract gluon TMDs. At this point, let us mention that back-to-back $J/\psi + Z$, which has been studied by ATLAS²⁵, could also be used to extract gluon TMDs. The arguments in favor of this observable are that it would probe them at higher scales ($Q \simeq m_Z$) and provide us with information on their evolution (see Ref. 29), that in disfavor is the very small expected yield³⁰. On the contrary, $J/\psi + W$ is likely contaminated by quark-induced contributions³¹. In addition, both $J/\psi + Z$ and $J/\psi + W$ are likely be contaminated by significant DPS contributions^{32,25}; a careful study of their suppression by the small-momentum-imbalance requirement would therefore be needed. The same proviso holds for quarkonium-pair production^{26,27,28}.

2. TMD formalism for photon-quarkonium production

In the TMD-factorization approach, the cross section for near back-to-back $Q + \gamma$ production is given by

$$d\sigma = \frac{1}{2s} \int dx_a dx_b d^2\mathbf{p}_{T,a} d^2\mathbf{p}_{T,b} \frac{d^3P_Q d^3P_\gamma}{16\pi^2 E_Q E_\gamma} \delta^4(p_a + p_b - P_Q - P_\gamma) \times \text{Tr} \{ \Phi_g(x_a, \mathbf{p}_{T,a}, \zeta, \mu) \Phi_g(x_b, \mathbf{p}_{T,b}, \zeta, \mu) |\mathcal{A}(g g \rightarrow Q \gamma)|^2 \}, \quad (4)$$

where $s = (P_A + P_B)^2$ is the total energy squared in the hadronic centre-of-mass frame. We stress that the q_T dependence of the cross section is completely factored out from the hard-scattering amplitude squared $|\mathcal{A}(g g \rightarrow Q \gamma)|^2$.




Fig. 2. Evaluation of the azimuthally-modulated q_T spectra $S_{q_T}^{(0)}$, $S_{q_T}^{(2)}$ and $S_{q_T}^{(4)}$ given in Eq. (8) for the process $pp \rightarrow Q\gamma X$, for $\sqrt{s} = 14$ TeV, $Q = 20$ GeV, $Y = 0$, $\theta = \pi/2$.

One then finds that

$$\frac{d\sigma}{dQdYd^2q_Td\Omega} = \frac{4\alpha_s^2\alpha^2e_Q^2|R_0(0)|^2}{3M_Q^2} \frac{Q^2 - M_Q^2}{sQ^3((\gamma^2 + 1)^2 - (\gamma^2 - 1)^2 \cos^2 \theta)} \times \left\{ F_1 C[f_1^g f_1^g] + F_3 C[w_3 f_1^g h_1^{\perp g} + (x_a \leftrightarrow x_b)] \cos 2\phi + F_4 C[w_4 h_1^{\perp g} h_1^{\perp g}] \cos 4\phi \right\}, \quad (5)$$

where Q and Y are, respectively, the invariant mass and the rapidity of the pair, $x_{a,b} = \exp[\pm Y] Q / \sqrt{s}$ and the solid angle $\Omega = (\theta, \phi)$ is measured in the Collins-Soper frame^b and

$$C[wfg] \equiv \int d^2p_{T,a} \int d^2p_{T,b} \delta^2(p_{T,a} + p_{T,b} - \mathbf{q}_T) w(p_{T,a}, p_{T,b}) f(x_1, p_{T,a}^2) g(x_2, p_{T,a}^2).$$

In Eq. (5), $R_0(0)$ is the quarkonium radial wave function at the origin in the position space and e_Q is the heavy-quark charge in units of the proton charge. The factors F_i are given by $F_1 = 1 + 2\gamma^2 + 9\gamma^4 + (6\gamma^4 - 2)\cos^2 \theta + (\gamma^2 - 1)^2 \cos^4 \theta$, $F_3 = 4\gamma^2 \sin^2 \theta$ and $F_4 = (\gamma^2 - 1)^2 \sin^4 \theta$ with $\gamma \equiv Q/M_Q$. The explicit expressions for the transverse weights are

$$w_3 = \frac{\mathbf{q}_T^2 \mathbf{p}_{bT}^2 - 2(\mathbf{q}_T \cdot \mathbf{p}_{bT})^2}{2M_p^2 \mathbf{q}_T^2}, w_4 = 2 \left[\frac{\mathbf{p}_{aT} \cdot \mathbf{p}_{bT}}{2M_p^2} - \frac{(\mathbf{p}_{aT} \cdot \mathbf{q}_T)(\mathbf{p}_{bT} \cdot \mathbf{q}_T)}{M_p^2 \mathbf{q}_T^2} \right]^2 - \frac{\mathbf{p}_{aT}^2 \mathbf{p}_{bT}^2}{4M_p^4}. \quad (6)$$

3. Extraction of f_1^g and $h_1^{\perp g}$

By measuring the following q_T -dependent observables,

$$S_{q_T}^{(n)} \equiv \frac{\int d\phi \cos(n\phi) \frac{d\sigma}{dQdYd^2q_Td\Omega}}{\int_0^{Q^2/4} d\mathbf{q}_T^2 \int d\phi \frac{d\sigma}{dQdYd^2q_Td\Omega}}, \quad (7)$$

^bThe Collins-Soper frame is such that the pair is at rest and the $\hat{x}\hat{z}$ -plane spanned by $(\mathbf{P}_A, \mathbf{P}_B)$ and the \hat{x} -axis set by their bisector³³.

with $n = 0, 2, 4$, one can single out the three terms in Eq. (5). We obtain

$$\begin{aligned} \mathcal{S}_{q_T}^{(0)} &= \frac{C[f_1^g f_1^g]}{\int d\mathbf{q}_T^2 C[f_1^g f_1^g]}, \\ \mathcal{S}_{q_T}^{(2)} &= \frac{F_3 C[w_3 f_1^g h_1^{\perp g} + (x_a \leftrightarrow x_b)]}{2F_1 \int d\mathbf{q}_T^2 C[f_1^g f_1^g]}, \\ \mathcal{S}_{q_T}^{(4)} &= \frac{F_4 C[w_4 h_1^{\perp g} h_1^{\perp g}]}{2F_1 \int d\mathbf{q}_T^2 C[f_1^g f_1^g]}. \end{aligned} \quad (8)$$

Fig. 2 shows predictions for $\mathcal{S}_{q_T}^{(0,2,4)}$ for the process $\Upsilon + \gamma$ with different Ansätze for the TMD distributions. We find that the size of $\mathcal{S}_{q_T}^{(0)}$ should be sufficient to allow for an extraction of f_1^g as a function of q_T . $\mathcal{S}_{q_T}^{(2)}$ and $\mathcal{S}_{q_T}^{(4)}$ are small and one would need to integrate them over \mathbf{q}_T^2 , (up to $Q^2/4$), to look for an experimental evidence of a nonzero $h_1^{\perp g}$.

4. Conclusion

We claim that a first experimental determination of the polarized and unpolarized gluon TMD distributions for x on the order of 10^{-3} can be obtained from the analyses of azimuthal asymmetries and transverse-momentum spectra in $p p \rightarrow J/\psi(\Upsilon) \gamma X$ at the LHC. The yields are large enough to perform these analyses using already existing data at the center-of-mass energies $\sqrt{s} = 7$ and 8 TeV.

Acknowledgments

I would like to thank Wilco den Dunnen, Cristian Pisano and Marc Schlegel with whom I have collaborated on the topic presented here. This work was supported in part by the French CNRS, Grants No. PICS-06149 Torino-IPNO and No. PEPS4AFTER2.

References

1. J.C. Collins, *Foundations of Perturbative QCD*, (Cambridge University Press, Cambridge, 2011); S.M. Aybat, T.C. Rogers, *Phys. Rev. D* **83**, 114042 (2011); J.C. Collins, T.C. Rogers, *Phys. Rev. D* **87**, 034018 (2013); M.G. Echevarria, A. Idilbi, I. Scimemi, *JHEP* **1207**, 002 (2012).
2. D. Boer, S.J. Brodsky, P.J. Mulders and C. Pisano, *Phys. Rev. Lett.* **106**, 132001 (2011).
3. C. Pisano, *et al.*, *JHEP* **1310**, 024 (2013).
4. D. Boer, P.J. Mulders and C. Pisano, *Phys. Rev. D* **80**, 094017 (2009).
5. T.C. Rogers and P.J. Mulders, *Phys. Rev. D* **81**, 094006 (2010).
6. J.W. Qiu, M. Schlegel and W. Vogelsang, *Phys. Rev. Lett.* **107**, 062001 (2011).
7. P.J. Mulders and J. Rodrigues, *Phys. Rev. D* **63**, 094021 (2001).
8. D. Boer, *et al.*, *Phys. Rev. Lett.* **108**, 032002 (2012).
9. D. Boer, W.J. den Dunnen, C. Pisano, M. Schlegel, *Phys. Rev. Lett.* **111**, 032002 (2013).
10. D. Boer and C. Pisano, *Phys. Rev. D* **86** (2012) 094007
11. J. P. Ma, J. X. Wang and S. Zhao, *Phys. Rev. D* **88** (2013) 014027
12. R. Aaij *et al.* [LHCb Collaboration], arXiv:1409.3612 [hep-ex].
13. J.P. Ma, J.X. Wang and S. Zhao, *Phys. Lett. B* **737** (2014) 103

14. S.J. Brodsky, F. Fleuret, C. Hadjidakis, J.P. Lansberg, *Phys. Rept.* **522** (2013) 239
15. J.P. Lansberg, S.J. Brodsky, F. Fleuret, C. Hadjidakis, *Few Body Syst.* **53** (2012) 11
16. J. P. Lansberg, *et al.* *EPJ Web of Conf.* **85**, 02038 (2015)
17. L. Massacrier, *et al.* arXiv:1502.00984 [nucl-ex].
18. W. J. den Dunnen, J. P. Lansberg, C. Pisano, M. Schlegel, *Phys. Rev. Lett.* **112** (2014) 212001
19. P. Mathews, K. Sridhar and R. Basu, *Phys. Rev. D* **60**, 014009 (1999).
20. N. Brambilla, *et al.* *Eur. Phys. J. C* **71** (2011) 1534
21. J. P. Lansberg, *Eur. Phys. J. C* **61** (2009) 693
22. J. P. Lansberg, *Int. J. Mod. Phys. A* **21** (2006) 3857
23. J. P. Lansberg, *Phys. Lett. B* **679** (2009) 340
24. D. Boer and C. Pisano, arXiv:1412.5556 [hep-ph].
25. G. Aad *et al.* [ATLAS Collaboration], arXiv:1412.6428 [hep-ex].
26. J. P. Lansberg and H. S. Shao, *Phys. Rev. Lett.* **111** (2013) 122001
27. G. P. Zhang, *Phys. Rev. D* **90** (2014) 9, 094011
28. J. P. Lansberg and H. S. Shao, arXiv:1410.8822 [hep-ph].
29. D. Boer and W. J. den Dunnen, *Nucl. Phys. B* **886** (2014) 421
30. B. Gong, J. P. Lansberg, C. Lorce and J. Wang, *JHEP* **1303** (2013) 115
31. J. P. Lansberg, C. Lorce, *Phys. Lett. B* **726** (2013) 218; erratum *Ibid.* **738** (2014) 529
32. G. Aad *et al.* [ATLAS Collaboration], *JHEP* **1404** (2014) 172
33. J.C. Collins and D.E. Soper, *Phys. Rev. D* **16**, 2219 (1977).